Securing the Health of Disadvantaged Women A Critical Investigation of Tobacco-Control Policy Effects

on Women Worldwide

Roland S. Moore, PhD, Deborah L. McLellan, MHS, MA, John A. Tauras, PhD, Pebbles Fagan, PhD, MPH

Introduction

obacco, directly or indirectly, affects almost every aspect of the lives of disadvantaged women and girls. For example, in many low- and middle-income countries such as Cambodia, China, the Philippines, and Viet Nam, 1.2 smoking is high among men and low among women. The diversion of income toward tobacco by male smokers in households contributes to malnutrition and secondhand smoke exposure among women and children. Tobacco accounts for 2.8% of the average household expenditures in poor countries like Bangladesh and 15% of total household expenditures among the lowest-income groups in Indonesia. Because of the high tobacco use rates among men, women in low-income countries are more likely to care for partners suffering from tobacco-related illnesses.

Among women working in service and manual labor positions,⁵ secondhand smoke exposure is high, and low-income women who manufacture tobacco in places like Southeast Asia and Brazil suffer from green tobacco sickness caused by transdermal exposure to tobacco.⁸ Further, with changes in social norms and increased female autonomy, tobacco use is increasing among women in low- and middle-income countries.⁹ This growing epidemic, which affects every aspect of the lives of women and girls, will ultimately increase lung cancer, other tobacco-related illnesses, second-hand smoke exposure, malnutrition, family economic disadvantage, and care-giving burden, as well as threaten food and financial security.

Global efforts to reduce the public health consequences of smoking among disadvantaged women call for the careful examination of how evidence-based interventions not only reduce tobacco use and exposure, but also affect the social context of their lives. Through the Framework Convention on Tobacco Con-

From the Pacific Institute for Research and Evaluation, Prevention Research Center (Moore), Berkeley, California; The Heller School for Social Policy and Management, Brandeis University (McLellan), Waltham, Massachusetts; the University of Chicago (Tauras), Chicago, Illinois; and the Division of Cancer Control and Population Sciences, National Cancer Institute, NIH (Fagan), Bethesda, Maryland

Address correspondence and reprint requests to: Roland S. Moore, PhD, Pacific Institute for Research and Evaluation, Prevention Research Center, 1995 University Avenue, Suite 450, Berkeley CA 94704. E-mail: roland@prev.org.

trol (FCTC), the WHO recently called for the global implementation of evidence-based tobacco policies, which can potentially reduce the global tobacco epidemic. In 2004, the Tobacco Research Network on Disparities (TReND; tobaccodisparities.org), co-funded by the National Cancer Institute and the American Legacy Foundation, launched the Low Socioeconomic Status Women and Girls Project to critically examine the full spectrum of the effects of evidence-based policies and programs on the health and well-being of disadvantaged women and girls.

During Phase I of this project, TReND convened a meeting of experts in 2005; published a special journal supplement¹⁰ in 2006; and published a report in 2008. 11 Phase I of this project concluded that low-SES women were less likely to have smoke-free homes^{4,5} than more-advantaged women, and worksite bans were not associated with quit attempts among low-SES women.⁶ Further, studies showed that gender and power dynamics play a key role in the enforcement of policies. 12,13 For example, in a study of randomly selected bars covered by smoke-free workplace laws, female bartenders were more likely to be exposed to tobacco smoke than their male counterparts. 13 Collectively, the studies showed that not all policies result in the intended effects, and there is a need to continue to examine the consequences of policies on populations with high rates of smoking, low rates of quitting, and at increased risk for tobacco-related diseases.

Theoretically, there are several reasons why unintended consequences may occur. One is that policies disrupt highly complex systems, in which reactions are not always predictable. Another reason is that psychological reactance (as in the case of educational campaigns or warning labels on alcohol or cigarettes) may produce the opposite (boomerang) effect from what is intended. Those addicted to nicotine may change brands or consumption patterns in order to restore intake levels altered by tobacco-control policies: Evidence indicates that young adults smoke higher-tar cigarettes in response to a cigarette tax hike.

Although some might conclude from the foregoing examples that unintended policy consequences are inevitable, ¹⁶ other research is more hopeful. Theory-based modeling of potential side effects of policy interventions in order to forestall unwanted outcomes

may help reduce such unintended consequences.¹⁷ Similarly, other scholars advocate for simulation modeling and complex systems thinking to avoid unwanted policy repercussions.^{18,19} Smoke-free workplace policies have negative unintended consequences including exposure to secondhand smoke for people entering and leaving work; policy improvements to ameliorate such problems have been implemented, such as extending no-smoking zones to 25 feet or more from building entrances.²⁰

In a variety of health policy domains, research has documented unintended consequences, especially for individuals with fewer resources. For example, restructuring Medicare drug benefit policies for fiscal efficiency increased emergency room visits and worsened physiological symptoms, especially for patients from neighborhoods characterized by low SES.²¹ In the realm of tobacco control, because regional smoke-free policies tend to be enacted earlier in higher SES communities, geographic and economic disparities in tobacco-control deployment constitute yet another way in which low-SES workers are exposed to greater amounts of secondhand smoke.²²

Restrictive marketing and promotion laws in the interest of health may inspire ever more creative workarounds by affected industries; one study²³ identified ways in which the tobacco industry reached out to youth in spite of laws designed to prevent them from doing so. Congruent with an analysis of internal tobacco industry documents in this supplement,²⁴ another study²⁵ showed how activists negotiating with the tobacco industry over warning labels inadvertently enhanced industry legitimacy.

These and other studies that examine how policies inadvertently cause harm and facilitate other healthful conditions or behaviors prompted the development of the second phase of the Low Socioeconomic Status Women and Girls Project: Unintended Consequences of Tobacco Policies on Low Socioeconomic Status Women and Girls. In 2007, TReND issued a global call for abstracts. Over 40 abstracts were submitted; papers meeting the guidelines were internally reviewed then peer-reviewed, and nine 24,26-33 are being published in this supplement to the American Journal of Preventive Medicine.

The guest editors required that all papers directly address: (1) a population of women and/or girls that clearly can be identified as being of low SES (typically but not always measured by such indicators as relative income, education, and occupational levels); (2) inclusion of an assessment of tobacco-control policy (such as tax increases; smoking restrictions in the home, car, or workplace; advertising restrictions); and (3) unintended consequences of tobacco-control policies. This journal supplement challenged authors to note unintended consequences from policies that could be either positive/helpful for low-SES females or negative/harmful

for them, across a variety of domains. In addition to the nine peer-reviewed papers, the supplement also features three commentaries, authored by Cheryl Healton, Donna Vallone, and Julie Cartwright³⁴; Hilary Graham³⁵; and Gloria Eldridge and Karen Cropsey.³⁶

Summary of Papers

The nine papers in this supplement 24,26-33 include literature reviews; qualitative, mixed-methods; and quantitative analyses of data collected in the U.S. and China. The Greaves and Hemsing²⁶ review suggests that low-income women living in high-density areas may have limited access to safe outdoor spaces or face child care issues if they have to go outside to smoke; that policies can contribute to the stigmatization of mothers who smoke, leading to women not seeking smokingcessation assistance; that women in service jobs are less likely to be protected by smoke-free policies because of a lack of enforcement or unequal distribution of policies; that smoking can increase in homes as a result of workplace and public policies; and that "smoking breaks" increase camaraderie among smokers. Similarly, Burgess et al.²⁷ suggest that low-SES groups are more likely to experience and internalize stigmatization compared to more-advantaged groups, and programs that are designed to reduce secondhand smoke exposure among children may stigmatize mothers as "bad mothers" and result in other unintended coping responses.

Several qualitative and mixed-methods studies captured helpful and harmful unintended effects of policies. Moore and colleagues³⁰ found that women who smoked outside smoke-free bars in California felt unsafe on the streets and had to deal with negative public perceptions and threats to safety in the rough neighborhoods in which some of these bars are located. Similar to Greaves and Hemsing, 26 Moore et al. 30 found that female patrons felt that smoking outside the bar provided an opportunity for social networking and solidarity. Yao et al. 33 examined the effects of workplace policies on pregnant women in Chengdu, China, and found that smoke-free policies at work or on public transit displaced men's smoking to home environments, increasing low-income women's exposure to secondhand smoke. These women also reported that their husbands were subject to greater pressure at work because of these policies and this pressure affected family harmony.

Balbach and Campbell²⁴ suggest that not acknowledging the economic effects of increased excise taxes on poor smokers could create an opportunity for the tobacco industry to form partnerships with groups representing low-income women. Working through the Tobacco Institute Labor Management Committee, the tobacco industry was successful at getting the Coalition of Labor Union Women to take a position on the

unfairness of excise taxes in exchange for financial incentives. Balbach and Campbell suggest that advocates and others need to be mindful of the opportunities policy proposals present to the tobacco industry.

Quantitative studies in this supplement focused on the harmful effects of smoke-free home and work policies, voucher, and tax policies. Fang and Rizzo²⁸ found that women who did not have a regular job and were less educated were more likely to start smoking during the time period when China began issuing inexpensive vouchers for the purchase of cigarettes. These vouchers were implemented from 1960 to 1980 as a means to control tobacco consumption and allow the Chinese government to use scarce resources on the daily necessities instead of cigarettes. Tong and colleagues³¹ reported that although Chinese and Korean-American women in California reported similar rates of smoke-free homes and work policies, lower-educated women were considerably more likely to be exposed to secondhand smoke than higher-educated women. These differences may be due to lack of control over the enforcement of policies and the empowerment of women in the home environment.

Hospitals in the U.S. have smoke-free environments that can affect the behaviors of mothers following birth. Paul and colleagues³² found that newborns of mothers who smoked had shorter mean length of stay than nonsmoking mothers. The more women smoked, the shorter the length of stay for newborns in Pennsylvania hospitals. Sarna et al.²⁹ found that nurses who did not smoke were more likely to miss breaks than nurses who were smokers. However, smoking status was not associated with missed breaks when comparing educational status; lower-educated nurses (LPNs) were more likely than more-educated nurses (RNs) to miss a break irrespective of smoking status.

Directions for Securing the Health and Well-Being of Disadvantaged Women

The devastating social and economic consequences of tobacco call for immediate global and concerted efforts to secure the health and well-being of disadvantaged women and girls. The results from these studies suggest that there are both harmful and beneficial effects of tobacco policies on low-SES women and girls. Stigmatization of mothers, lack of enforcement and ability to negotiate smoking policy in the home or at work, the neglect of tobacco researchers and advocates to carefully consider the needs of low-SES women and groups representing these women, safety issues, and the perception that work and public policies "shift" smoking to homes were consistent themes in one or more papers. In addition, several papers found one positive effect work breaks and smoking outside resulted in increased camaraderie and solidarity. These studies collectively

send a strong message that we need to move beyond the traditional narrow focus of examining how tobacco policies affect smoking, quitting, and secondhand smoke exposure. Studies confirm that passing policies alone is an important but not sufficient step. There is an urgent need to develop broader frameworks that are gender sensitive, ¹² recognize the complexity and interactions of societal systems, and move us toward greater progress in reducing inequities in tobacco use, exposure, and the consequences experienced by low-SES women and girls.

Recognition of the complexity of the problem prompts us to take advantage of extraordinary opportunities in the coming years. In countries that have ratified the Framework Convention on Tobacco Control, a socio-gender-sensitive agenda obligates us to examine positive and negative consequences of increases in taxes, clean indoor air laws, youth access laws, warning labels, marketing and production restrictions, prevention of illicit sales and trade, and voluntary home policies on disadvantaged women and girls. An even broader agenda prompts us to examine tobacco policy interactions with social policies such as the Children's Health Insurance Program in the U.S. The Children's Health Insurance Program Reauthorization Act of 2009, which expands health and dental coverage to 4 million uninsured low-income children and pregnant women,³⁷ provides not only the opportunity to evaluate the intended and unintended consequences of the \$0.62 tax increase on disadvantaged women, but more importantly, allows us to examine the interactive effects of insurance coverage and tobacco policies on the health and well-being of disadvantaged women and girls. The 2009 Lilly Ledbetter Fair Pay Act that addresses pay equity and discriminatory action by employers for each discriminatory paycheck,³⁸ also presents an extraordinary opportunity to redress how pay equity helps or harms the health outcomes and social circumstances of disadvantaged women and girls. These and examples from other countries are numerous—careful and thoughtful socio-gender-specific agenda setting is necessary so that the different needs of disadvantaged women and the pros and cons of policies be considered at all stages of policy and program development in legislative or voluntary systems.

As the studies in this supplement suggest, the formulation of gender- and SES-sensitive policy must occur across social, political, economic, healthcare, housing, education, justice, food security, agriculture, commerce, and other systems whose policies—and their interactions with tobacco policies—influence the health and well-being of disadvantaged women. Personal, political, and fiscal commitment and innovation are needed to improve health and health equity across the life course of women and girls. Research innovation and bold practical solutions, and the will and resilience

of individuals from all socioeconomic classes are also necessary.

Developing practical solutions will require policy-makers to hold meaningful discussions with low-SES women and organizations that represent them on complimentary programs that alleviate any detrimental effects. More than ever, research on policy repercussions, especially among disadvantaged women, is needed in order to build the collective skills and capacity to implement and promote appropriate policies and complimentary programs that secure the health and well-being of the families around the globe.

The authors gratefully acknowledge the valuable contributions of TReND network members in the internal review process. They also thank Allison Rose and Julie Murphy for bibliographic assistance.

No financial disclosures were reported by the authors of this paper.

References

- Bloch M, Althabe F, Onyamboko M, et al. Tobacco use and secondhand smoke exposure during pregnancy: an investigative survey of women in 9 developing nations. Am J Public Health 2008;98:1833–40.
- WHO. Tobacco. In: World Health Organization Regional Office for the Western Pacific, 2009.
- Semba RD, Kalm LM, de Pee S, Ricks MO, Sari M, Bloem MW. Paternal smoking is associated with increased risk of child malnutrition among poor urban families in Indonesia. Publ Health Nutr 2007;10:7–15.
- Shavers VL, Fagan P, Alexander LA, Clayton R, Doucet J, Baezconde-Garbanati L. Workplace and home smoking restrictions and racial/ethnic variation in the prevalence and intensity of current cigarette smoking among women by poverty status, TUS-CPS 1998–1999 and 2001–2002. J Epidemiol Community Health 2006;60(2S):34–43.
- Shopland DR, Anderson CM, Burns DM. Association between home smoking restrictions and changes in smoking behaviour among employed women. J Epidemiol Community Health 2006;60(2S):44–50.
- Efroymson D, Ahmed S, Townsend J, et al. Hungry for tobacco: an analysis
 of the economic impact of tobacco consumption on the poor in Bangladesh. Tob Control 2001;10:212–7.
- 7. Barraclough S. Women and tobacco in Indonesia. Tob Control
- Arcury TA, Quandt SA, Preisser JS, Bernert JT, Norton D, Wang J. High levels of transdermal nicotine exposure produce green tobacco sickness in Latino farmworkers. Nicotine Tob Res 2003;5:315–21.
- WHO. WHO report on the global tobacco epidemic, 2008: The MPOWER package. Geneva: WHO, 2008.
- Tobacco Control Policy and Low Socioeconomic Status Women and Girls. I Epidemiol Community Health 2006;60(28):1–65.
- TReND: Low SES Women and Girls Project (Phase I): The Effects
 of Tobacco Control Policies on Low SES Women and Girls.
 cancercontrol.cancer.gov/tcrb/ses_women-girls_project/index.html.
- Greaves L, Jategaonkar N. Tobacco policies and vulnerable girls and women: toward a framework for gender sensitive policy development. J Epidemiol Community Health 2006;60(2S):57–65.
- Moore RS, Lee JP, Antin TMJ, Martin SE. Tobacco free workplace policies and low socioeconomic status female bartenders in San Francisco. J Epidemiol Community Health 2006;60(2S):51–56.

- Ringold DJ. Boomerang effects in response to public health interventions: some unintended consequences in the alcoholic beverage market. J Consumer Policy 2002;25:27–63.
- Evans WN, Farrelly MC. The compensating behavior of smokers: taxes, tar, and nicotine. Rand J Econ 1998;29:578–95.
- Roots RI. When laws backfire: unintended consequences of public policy. Am Behav Sci 2004;47:1376–94.
- Rosenberg BJ, Barbeau EM, Moure-Eraso R, Levenstein C. The work environment impact assessment: a methodologic framework for evaluating health-based interventions. Am J Ind Med 2001;39:218–26.
- Sterman JD. Learning from evidence in a complex world. Am J Public Health 2006;96:505–14.
- Thomson R, Murtagh M, Khaw FM. Tensions in public health policy: patient engagement, evidence-based public health and health inequalities. Qual Saf Health Care 2005;14:398–400.
- Parry O, Platt S, Thomson C. Out of sight, out of mind: workplace smoking bans and the relocation of smoking at work. Health Promotion Intl 2000;15:125–33.
- Hsu J, Price M, Huang J, et al. Unintended consequences of caps on Medicare drug benefits. N Engl J Med 2006;354:2349–59.
- Skeer M, George S, Hamilton WL, Cheng DM, Siegel M. Town-level characteristics and smoking policy adoption in Massachusetts: are local restaurant smoking regulations fostering disparities in health protection? Am J Public Health 2004;94:286–92.
- Hoek J. Tobacco promotion restrictions: ironies and unintended consequences. J Business Res 2004;57:1250-7.
- Balbach ED, Campbell RB. Union women, the tobacco industry, and excise taxes: a lesson in unintended consequences. Am J Prev Med 2009; 37(2S):S121–S125.
- Wander N, Malone RE. Making big tobacco give in: you lose, they win. Am J Public Health 2006;96:2048–54.
- Greaves L, Hemsing N. Sex, gender, and secondhand smoke policies: implications for disadvantaged women. Am J Prev Med 2009;37(2S): S131–S137.
- Burgess DJ, Fu SS, van Ryn M. Potential unintended consequences of tobacco-control policies on mothers who smoke: a review of the literature. Am J Prev Med 2009;37(2S):S151–S158.
- Fang H, Rizzo JA. Did cigarette vouchers increase female smokers in China?
 Am J Prev Med 2009;37(2S):S126–S130.
- Sarna L, Bialous SA, Wells M, Kotlerman J, Froelicher ES, Wewers ME. Do you need to smoke to get a break? Smoking status and missed work breaks among staff nurses. Am J Prev Med 2009;37(2S):S165–S171.
- 30. Moore RS, Annechino R, Lee JP. Unintended consequences of smoke-free bar policies for low-SES women in three California counties. Am J Prev Med 2009;37(2S):S138–S143.
- 31. Tong EK, Tang H, Tsoh J, Wong C, Chen MS. Jr. Smoke-free policies among Asian-American women: comparisons by education status. Am J Prev Med 2009;37(2S):S144–S150.
- Paul IM, Lehman EB, Widome R. Maternal tobacco use and shorter newborn nursery stays. Am J Prev Med 2009;37(28):S172–S178.
- 33. Yao T, Lee AH, Mao Z. Potential unintended consequences of smoke-free policies in public places on pregnant women in China. Am J Prev Med 2009;37(2S):S159–S164.
- Healton CG, Vallone D, Cartwright J. Unintended consequences of tobacco policies: implications for public health practice. Am J Prev Med 2009;37(2S):S181–S182.
- Graham H. Why social disparities matter for tobacco-control policy. Am J Prev Med 2009;37(2S):S183–S184.
- 36. Eldridge GC, Cropsey KL. Smoking bans and restrictions in U.S. prisons and jails: consequences for incarcerated women. Am J Prev Med 2009;37(2S):S179-S180.
- 37. Cuttler L, Kenney GM. Update on the State Children's Health Insurance Program. Arch Pediatr Adolesc Med 2009;163:103–7.
- Stolberg SG. Obama signs equal-pay legislation. New York Times 2009 January 29, 2009; Sect. A1.